

Management of Chronic Prostatitis/ Chronic Pelvic Pain Syndrome

A Systematic Review and Network Meta-analysis

Thunyarat Anothaisintawee, MD

John Attia, MD, PhD, FRCPC

J. Curtis Nickel, MD, FRCSC

Sangsuree Thammakraisorn, MD

Pawin Numthavaj, MD

Mark McEvoy, MSc

Ammarin Thakkinian, PhD

PROSTATITIS IS A COMMON condition, with an estimated prevalence in the community of about 9%,¹ and accounts for nearly 2 million ambulatory care encounters annually in the United States.² However, prostatitis represents a heterogeneous mix of conditions, including acute prostatitis, chronic bacterial prostatitis, and asymptomatic inflammatory prostatitis. Chronic prostatitis/chronic pelvic pain syndrome (CP/CPPS), which accounts for 90% to 95% of cases,^{3,4} is a clinical entity defined as urologic pain or discomfort in the pelvic region, associated with urinary symptoms and/or sexual dysfunction, lasting for at least 3 of the previous 6 months. Chronic prostatitis/chronic pelvic pain syndrome is a diagnosis of exclusion that can be made after ruling out active urethritis, urogenital cancer, urinary tract disease, urethral stricture, or neurological disease affecting the bladder. Symptoms of CP/CPPS can diminish quality of life and impair physical and psychological function.⁵

The etiology of CP/CPPS is uncertain but may include inflammatory or non-inflammatory etiologies.⁶⁻⁸ An inciting agent may cause inflammation or neurological damage in or around the prostate

Context Chronic prostatitis/chronic pelvic pain syndrome (CP/CPPS) is common, but trial evidence is conflicting and therapeutic options are controversial.

Objective To conduct a systematic review and network meta-analysis comparing mean symptom scores and treatment response among α -blockers, antibiotics, anti-inflammatory drugs, other active drugs (phytotherapy, glycosaminoglycans, finasteride, and neuromodulators), and placebo.

Data Sources We searched MEDLINE from 1949 and EMBASE from 1974 to November 16, 2010, using the PubMed and Ovid search engines.

Study Selection Randomized controlled trials comparing drug treatments in CP/CPPS patients.

Data Extraction Two reviewers independently extracted mean symptom scores, quality-of-life measures, and response to treatment between treatment groups. Standardized mean difference and random-effects methods were applied for pooling continuous and dichotomous outcomes, respectively. A longitudinal mixed regression model was used for network meta-analysis to indirectly compare treatment effects.

Data Synthesis Twenty-three of 262 studies identified were eligible. Compared with placebo, α -blockers were associated with significant improvement in symptoms with standardized mean differences in total symptom, pain, voiding, and quality-of-life scores of -1.7 (95% confidence interval [CI], -2.8 to -0.6), -1.1 (95% CI, -1.8 to -0.3), -1.4 (95% CI, -2.3 to -0.5), and -1.0 (95% CI, -1.8 to -0.2), respectively. Patients receiving α -blockers or anti-inflammatory medications had a higher chance of favorable response compared with placebo, with pooled RRs of 1.6 (95% CI, 1.1-2.3) and 1.8 (95% CI, 1.2-2.6), respectively. Contour-enhanced funnel plots suggested the presence of publication bias for smaller studies of α -blocker therapies. The network meta-analysis suggested benefits of antibiotics in decreasing total symptom scores (-9.8 ; 95% CI, -15.1 to -4.6), pain scores (-4.4 ; 95% CI, -7.0 to -1.9), voiding scores (-2.8 ; 95% CI, -4.1 to -1.6), and quality-of-life scores (-1.9 ; 95% CI, -3.6 to -0.2) compared with placebo. Combining α -blockers and antibiotics yielded the greatest benefits compared with placebo, with corresponding decreases of -13.8 (95% CI, -17.5 to -10.2) for total symptom scores, -5.7 (95% CI, -7.8 to -3.6) for pain scores, -3.7 (95% CI, -5.2 to -2.1) for voiding, and -2.8 (95% CI, -4.7 to -0.9) for quality-of-life scores.

Conclusions α -Blockers, antibiotics, and combinations of these therapies appear to achieve the greatest improvement in clinical symptom scores compared with placebo. Anti-inflammatory therapies have a lesser but measurable benefit on selected outcomes. However, beneficial effects of α -blockers may be overestimated because of publication bias.

JAMA. 2011;305(1):78-86

www.jama.com

and lead to pelvic floor neuromuscular and/or neuropathic pain. Predisposing factors for CP/CPPS may include heredity, infection, voiding abnormalities, hor-

mone imbalance, intraprostatic reflux, immunological or allergic triggers, or psychological traits. A wide variety of therapies including α -blockers, antibiotics,

 CME available online at www.jamaarchivescme.com and questions on p 105.

Author Affiliations are listed at the end of this article. **Corresponding Author:** Ammarin Thakkinian, PhD, Section for Clinical Epidemiology and Biostatistics, Ramathibodi Hospital, Rama VI Rd, Rachatevi, Bangkok, Thailand 10400 (raatk@mahidol.ac.th).

Clinical Review Section Editor: Mary McGrae McDermott, MD, Contributing Editor. We encourage authors to submit papers for consideration as a Clinical Review. Please contact Mary McGrae McDermott, MD, at mdm608@northwestern.edu.

anti-inflammatory medications, and other agents (eg, finasteride, phytotherapy, and gabapentinoids) are routinely used. However, the efficacy of these treatments is controversial,⁹⁻¹⁵ partly because many clinical trials testing these therapies have been small, with little statistical power to detect treatment effects.

To date, only 1 systematic review⁶ and 1 meta-analysis¹⁶ of α -blockers vs placebo of which we are aware have been performed for treatment of CP/CPPS. We therefore performed a systematic review and network meta-analysis mapping all treatment regimens, with 2 aims. First, we compared total symptom, pain, voiding, and quality-of-life scores at the end of therapy with α -blockers (the most commonly evaluated therapy for CP/CPPS), other active drugs, or placebo. Second, we compared rates of responses to therapies available for treating CP/CPPS.

METHODS

Search Strategy

We searched the MEDLINE and EMBASE databases for relevant studies published in English from 1949 (for MEDLINE) or 1974 (for EMBASE) through November 16, 2010. Search terms and strategies for each database are described in the eAppendix (available at <http://www.jama.com>). Reference lists of included trials and the previous systematic reviews^{6,16} were explored.

Selection of Studies

Identified studies were selected based on title and abstract by 2 independent authors (T.A. and S.T.). Full articles were retrieved if a decision could not be made based on the abstracts. Agreement between the 2 reviewers was measured using the κ statistic. Disagreement was resolved by consensus and by discussion with a third party (J.C.N. or A.T.).

Inclusion Criteria

Randomized controlled trials that were published in English were selected if they met the following criteria: (1) Participants met criteria for CP/CPPS categories IIIA or IIIB according to the National Institutes of Health classification.⁴ (2) The

study compared any pair of the following interventions: α -blockers, antibiotics, steroidal and nonsteroidal anti-inflammatory drugs, finasteride, glycosaminoglycans, phytotherapy, gabapentinoids, and placebo. (3) The study measured any of the following outcomes: pain scores, voiding scores, quality-of-life scores, and total symptom scores. The total symptom score is a summation of pain, voiding, and quality-of-life scores. (4) The full article could be retrieved and had sufficient data for extraction, including number of patients, means and standard deviations of continuous outcomes in each group, and/or numbers of patients per group for dichotomous outcomes.

Data Extraction

Two authors (T.A. and S.T.) independently extracted data using a standardized extraction form. Disagreement was resolved by discussion or consensus with a third party (A.T.). Missing information was sought by contacting the corresponding authors of the studies.

Risk of Bias Assessment

Two authors (T.A. and P.N.) independently assessed risk of bias of each study using an established tool.¹⁷ Six domains were assessed: sequence generation, allocation concealment, blinding, incomplete outcome data, selective outcome reporting, and other sources of bias. Disagreement between 2 authors was resolved by consensus and discussion. Levels of agreement for each domain and the overall domains were assessed using the κ statistic.

Outcomes

The outcomes of interests were total symptom, pain, voiding, and quality-of-life scores and response rates as defined in the original articles. The following tools were used in these assessments: (1) The National Institutes of Health Chronic Prostatitis Symptom Index (NIH-CPSI) consists of 3 domains (ie, pain, voiding, and quality of life). The total scores range from 0 to 43.¹⁸ (2) The International Prostate Symptom Score (IPSS) questionnaire¹⁹ consists of 3 domains (ie,

pain, voiding, and quality of life), with combined scores ranging from 0 to 51. (3) The Prostatitis Symptom Score Index (PSSI) measures only pain scores, with a total score of 0 to 12. (4) Other pain and voiding questionnaires²⁰ consist of 7 and 5 items, respectively, with each item graded as 0 to 3.

For each measurement, scores closer to 0 reflect more favorable status. The minimal clinical significant difference for all these scales is 3 to 6 points.²¹⁻²⁴ For response to treatment, various definitions were used in the original studies; eg, 25%, 33%, or 50% decreases in the NIH-CPSI or 4 (clinically perceptible improvement) to 6 (clinically significant improvement) unit score decreases in the NIH-CPSI from baseline.

Statistical Analysis

For the direct meta-analysis, mean differences of continuous outcomes (ie, total symptom, pain, voiding, and quality-of-life scores) between treatment groups were estimated for each study and were pooled using a standardized mean difference (SMD) if the scales used for outcome measures differed. Otherwise, an unstandardized mean difference (USMD) was applied. Heterogeneity of the mean difference was assessed using Q and I^2 statistics. If heterogeneity was present or the degree of heterogeneity (I^2) was greater than 25%, the SMD was estimated using a random-effects model. Otherwise, a fixed-effects model was applied.

Relative risks (RRs) of response to treatment were estimated for each study. If there was evidence of heterogeneity, a random-effects model was used for pooling. Otherwise, the inverse-variance method was used. The source of heterogeneity was explored by fitting covariables (ie, mean age, duration of treatment, and baseline total symptom scores) one by one in the meta-regression. Publication bias was assessed using contour-enhanced funnel plots and the Egger test.^{25,26}

For indirect comparisons, network meta-analyses were applied to assess treatment effects for all possible treatment groups if summary data were available.²⁷⁻²⁹ Linear regression models weighted by inverse variance were applied to assess treat-

ment effects for continuous outcomes. Effects of study were included as covariates in the model. For response to treatment, summary data were expanded to individual patient-level data using the “expand” command in Stata. Treatment groups were considered in a mixed-effect hierarchical model with a log-link function using the “xtpoisson” command.²⁷ Pooled RRs and 95% confidence intervals (CIs) were estimated by exponential coefficients of treatments. All analyses were performed using Stata software, version 11.0.³⁰ Two-sided $P < .05$ was considered statistically significant except for the heterogeneity test, in which a 1-sided $P < .10$ was used.

RESULTS

Among 262 identified studies, 25 studies were eligible for inclusion (FIGURE). Two studies^{31,32} had insufficient data (ie, did not report means and standard de-

viations). For these studies, one corresponding author³¹ was contacted for additional information but did not respond, while the author of the other study³² could not be contacted. This left 23 studies^{9-15,21,22,24,33-45} with sufficient data for extraction. Agreement on study selection between the 2 reviewers was high at 98% ($\kappa = 0.91$; $P < .001$). Disagreements were present for 5 studies (selected by one reviewer but not the other) and included 2 duplicate reports, 1 non-randomized controlled trial, 1 protocol report, and 1 study with mixed CP/CPPS and bacterial prostatitis patients.

Twenty studies* compared outcomes between 1 active treatment and placebo (TABLE 1). These treatments were α -blockers in 7 studies,^{9,10,24,33-35,38} antibiotics in 2 studies,^{39,44} finasteride in 2 studies,^{37,40} anti-inflammatory drugs in

*References 9-15, 22, 24, 33-35, 37-41, 43-45.

4 studies,^{11,12,22,43} phytotherapy in 3 studies,¹³⁻¹⁵ glycosaminoglycan in 1 study,⁴¹ and pregabalin in 1 study.⁴⁵ Three studies^{21,36,42} had more than 2 treatment groups as follows: α -blockers plus antibiotics, α -blockers alone, and antibiotics alone in 2 studies^{36,42}; and α -blockers plus antibiotics, α -blockers alone, antibiotics alone, and placebo in 1 study.²¹ Treatment duration ranged from 4 to 52 weeks. Most studies used NIH-CPSI scores for measurement of outcomes. Mean age of participants ranged from 29.1 to 56.1 years.

eTable 1 reports the quality assessments of included studies. The agreement between 2 reviewers for each bias assessment domain ranged from 56% to 100% and the overall agreement was 91%. The highest quality was in the domain of selective outcome reports (95.7% low-risk) followed by blinding (87.4%). The lowest quality was in allocation concealment (adequate in 21.7%).


Direct Meta-analysis

Pooled Mean Scores. Eight studies compared mean scores between α -blockers and placebo. Among them, 4 studies^{10,33-35} reported mean scores at follow-up, while the remainder^{9,21,24,38} reported change in mean score. Among the latter, 2 studies^{9,21} reported information that allowed data simulation and pooling.^{9,10,21,33,34}

Three studies^{21,39,44} compared antibiotics with placebo and 3 studies¹³⁻¹⁵ compared phytotherapy with placebo.

Total Symptom Scores. Five studies^{9,10,21,33,34} (n = 568) comparing α -blockers and placebo were pooled (TABLE 2). Total symptom scores were assessed at the end of treatment, which ranged from 6 to 24 weeks. Mean differences and 95% CIs are shown in eFigure 1A. The total symptom score SMD in the α -blocker group vs the placebo group was -1.7 (95% CIs, -2.8 to -0.6), with high heterogeneity ($I^2 = 96.4\%$). This is equivalent to -5.5 (95% CI, -10.0 to -0.9) units on the NIH-CPSI or IPSS scales. Meta-regression did not identify the source of heterogeneity. Sensitivity analysis was performed by pooling the 4 studies^{9,21,33,34} with adequate sequence generation of randomization.

Figure. Study Selection



CP/CPPS indicates chronic prostatitis/chronic pelvic pain syndrome.

Table 1. Characteristics of Included Studies

Source	No. of Participants (Intervention)	Duration of Treatment, wk	Outcome Assessment Tool ^a	Age, Mean (SD)	Total Symptom Score, Mean (SD) ^b
α-Blocker vs placebo					
Nickel et al, ⁹ 2008	138 (Alfuzosin) 134 (Placebo)	12	NIH-CPSI	40.1 (1.4)	24.4 (0.7)
Tuçcu et al, ¹⁰ 2007	30 (Doxazosin) 30 (Placebo)	24	NIH-CPSI	29.1 (5.2)	23.0 (0.4)
Alexander et al, ²¹ 2004	49 (Tamsulosin) 49 (Placebo)	6	NIH-CPSI	44.6 (3.2)	24.8 (1.7)
Nickel et al, ²⁴ 2004	27 (Tamsulosin) 30 (Placebo)	6	NIH-CPSI	40.8 (21-56) ^c	26.3
Cheah et al, ³³ 2003	43 (Terazosin) 43 (Placebo)	14	NIH-CPSI	35.5 (20-50) ^c	26.2 (1.6)
Evlıyaoğlu and Burgut, ³⁴ 2002	30 (Doxazosin) 30 (Placebo)	12	IPSS, pain questionnaire	33.0 (28.5)	17.2 (1.0)
Gül et al, ³⁵ 2001	39 (Terazosine) 30 (Placebo)	12	PSSI	39.6	NR
Mehik et al, ³⁶ 2003	19 (Alfuzosin) 21 (Placebo)	24	NIH-CPSI	49.5	24.4
Antibiotic vs placebo					
Alexander et al, ²¹ 2004	49 (Ciprofloxacin) 49 (Placebo)	6	NIH-CPSI	44.6 (3.2)	24.8 (1.7)
Nickel et al, ³⁹ 2003	45 (Levofloxacin) 35 (Placebo)	6	NIH-CPSI	56.1 (36-78) ^c	23.0 (1.7)
Zhou et al, ⁴⁴ 2008	24 (Tetracycline hydrochloride) 24 (Placebo)	12	NIH-CPSI	NR	34.3 (1.2)
Finasteride vs placebo					
Leskinen et al, ³⁷ 1999	31 (Finasteride) 10 (Placebo)	52	IPSS, pain questionnaire	46.7 (32-61) ^c	NR
Nickel et al, ⁴⁰ 2004	33 (Finasteride) 31 (Placebo)	24	NIH-CPSI	44.4 (0.5)	21.3 (0.4)
Anti-inflammatory drugs vs placebo					
Goldmeier et al, ¹¹ 2005	10 (Zafirlukast) 7 (Placebo)	4	NIH-CPSI	35.9 (5.7)	NR
Nickel et al, ¹² 2003	49 (Rofecoxib) 59 (Placebo)	6	NIH-CPSI	46.8 (2.5)	21.8 (1.1)
Bates et al, ²² 2007	9 (Prednisolone) 12 (Placebo)	4	NIH-CPSI	40.8 (4.6)	24.3 (3.0)
Zhao et al, ⁴³ 2009	32 (Celecoxib) 32 (Placebo)	6	NIH-CPSI	NR	24.4 (1.4)
Phytotherapy vs placebo					
Elist, ¹³ 2006	30 (Pollen extract) 30 (Placebo)	24	Pain/voiding questionnaires	35.0 (20-55) ^c	
Shoskes et al, ¹⁴ 1999	15 (Bioflavonoid) 15 (Placebo)	4	NIH-CPSI	44.9 (5.4)	20.6 (2.1)
Wagenlehner et al, ¹⁵ 2009	70 (Pollen extract) 69 (Placebo)	12	NIH-CPSI	39.5 (8.1)	19.8 (5.2)
Glycosaminoglycan vs placebo					
Nickel et al, ⁴¹ 2005	51 (Pentosan polysulfate) 49 (Placebo)	16	NIH-CPSI	39.2 (21-59) ^c	26.5 (1.6)
Pregabalin vs placebo					
Pontari et al, ⁴⁵ 2010	217 (Pregabalin) 104 (Placebo)	6	NIH-CPSI	NR	26.1 (5.7)
Dual therapy vs monotherapy					
Alexander et al, ²¹ 2004	49 (Ciprofloxacin + tamsulosin) 49 (Tamsulosin) 49 (Ciprofloxacin) 49 (Placebo)	6	NIH-CPSI	44.6 (3.2)	24.8 (1.7)
Jeong et al, ³⁶ 2008	29 (Doxazosin + levofloxacin) 26 (Doxazosin) 26 (Levofloxacin)	6	NIH-CPSI	40.1 (23-60) ^c	23.1 (2.2)
Ye et al, ⁴² 2008	42 (Tamsulosin + levofloxacin) 42 (Tamsulosin) 21 (Levofloxacin)	12	NIH-CPSI	NR	27.6

Abbreviation: NR, data not reported.

^aInternational Prostate Symptom Score (IPSS) ranges from 0 to 51; National Institutes of Health Chronic Prostatitis Symptom Index (NIH-CPSI) ranges from 0 to 43; Prostatitis Symptom Score Index (PSSI) ranges from 0 to 12; and pain and voiding questionnaires range from 0 to 3.

^bMeasured at baseline, before receiving treatment.

^cRange is given instead of standard deviation.

In this analysis, the treatment effect was still significant, with a USMD of -4.0 (95% CI, -6.4 to -1.6). The Egger test indicated the presence of publication bias (coefficient = -9.3; SE, 2.5; P = .03) (eFigure 2A). Adjusting for publication bias using regression-based analy-

sis resulted in no evidence of treatment benefit (coefficient = 0.47; P = .39). Three studies^{21,39,44} (n = 215) were pooled to compare antibiotics with placebo (Table 2). The USMD was -5.8 (95% CI, -15.9 to 4.4). That is, the mean total symptom score in the antibiotic group was

5.8 units lower on the NIH-CPSI scale than in the placebo group, but this did not reach statistical significance (P = .26).

Pain Scores. Six studies^{9,10,21,33-35} (n = 637) compared mean pain scores between α -blockers and placebo (Table 2 and eFigure 1B). The SMD in the ran-

Table 2. Mean Symptom Scores at the End of Therapy According to Treatment

Source	Outcome Assessment Tool ^a	Active Treatment		Placebo	
		No. of Participants	Score, Mean (SD)	No. of Participants	Score, Mean (SD)
Total Symptom Score					
α-Blockers					
Nickel et al, ⁹ 2008	NIH-CPSI	138	16.7 (14.9)	134	18.6 (14.1)
Tuğcu et al, ¹⁰ 2007	NIH-CPSI	30	10.7 (1.3)	30	21.9 (1.2)
Alexander et al, ²¹ 2004	NIH-CPSI	45	20.2 (12.2)	45	21.6 (9.8)
Cheah et al, ³³ 2003	NIH-CPSI	43	10.8 (9.0)	43	17.0 (12.1)
Evliyaoğlu and Burgut, ³⁴ 2002	IPSS, pain questionnaire	30	10.5 (4.4)	30	16.2 (5.7)
SMD (95% CI)		-1.7 (-2.8 to -0.6)			
Antibiotics					
Alexander et al, ²¹ 2004	NIH-CPSI	42	18.0 (13.2)	45	21.6 (9.8)
Nickel et al, ³⁹ 2003	NIH-CPSI	45	19.0 (9.5)	35	18.4 (9.1)
Zhou et al, ⁴⁴ 2008	NIH-CPSI	24	17.1 (2.8)	24	31 (2.5)
USMD (95% CI)		-5.8 (-15.9 to 4.4)			
Pain Score					
α-Blockers					
Nickel et al, ⁹ 2008	NIH-CPSI	138	7.8 (7.6)	134	8.5 (7.6)
Tuğcu et al, ¹⁰ 2007	NIH-CPSI	30	4.7 (1.2)	30	8.6 (0.8)
Alexander et al, ²¹ 2004	NIH-CPSI	45	9.0 (7.1)	45	10.6 (5.8)
Cheah et al, ³³ 2003	NIH-CPSI	43	5.2 (5.7)	43	7.8 (6.7)
Evliyaoğlu and Burgut, ³⁴ 2002	Pain questionnaire	30	2.3 (1.1)	30	3.7 (1.3)
Gül et al, ³⁵ 2001	PSSI	39	6.3 (1.6)	30	8.8 (2.7)
SMD (95% CI)		-1.1 (-1.8 to -0.4)			
Antibiotics					
Alexander et al, ²¹ 2004	NIH-CPSI	42	8.7 (6.7)	45	10.6 (5.8)
Nickel et al, ³⁹ 2003	NIH-CPSI	45	8.9 (5.0)	35	7.6 (4.7)
Zhou et al, ⁴⁴ 2008	NIH-CPSI	24	7.1 (1.0)	24	14.5 (2.0)
USMD (95% CI)		-2.7 (-8.7 to 3.2)			
Phytotherapy					
Elist, ¹³ 2006	Pain questionnaire	30	2.8 (3.3)	28	4.9 (3.8)
Shoskes et al, ¹⁴ 1999	NIH-CPSI	15	6.2 (3.9)	13	9.0 (3.2)
Wagenlehner et al, ¹⁵ 2009	NIH-CPSI	68	5.5 (4.9)	68	7.3 (4.9)
SMD (95% CI)		-0.5 (-0.7 to -0.2)			
Voiding Score					
α-Blockers					
Nickel et al, ⁹ 2008	NIH-CPSI	138	3.3 (4.0)	134	3.9 (4.1)
Tuğcu et al, ¹⁰ 2007	NIH-CPSI	30	2.2 (0.8)	30	6.4 (1.0)
Alexander et al, ²¹ 2004	NIH-CPSI	45	4.2 (4.0)	45	10.6 (5.8)
Cheah et al, ³³ 2003	NIH-CPSI	43	2.0 (2.8)	43	3.6 (3.6)
Evliyaoğlu and Burgut, ³⁴ 2002	IPSS	30	5.9 (2.5)	30	8.8 (3.6)
SMD (95% CI)		-1.4 (-2.3 to -0.5)			
Antibiotics					
Alexander et al, ²¹ 2004	NIH-CPSI	42	3.5 (3.8)	45	10.6 (5.8)
Nickel et al, ³⁹ 2003	NIH-CPSI	45	4.2 (3.0)	35	4.1 (2.8)
Zhou et al, ⁴⁴ 2008	NIH-CPSI	24	5 (0.8)	24	8 (0.5)
USMD (95% CI)		-3.2 (-6.1 to -0.3)			
Phytotherapy					
Elist, ¹³ 2006	Pain questionnaire	30	1.4 (2.1)	28	2.8 (2.8)
Shoskes et al, ¹⁴ 1999	NIH-CPSI	15	1.5 (1.9)	13	3.0 (2.7)
Wagenlehner et al, ¹⁵ 2009	NIH-CPSI	68	1.8 (2.9)	69	2.6 (3.1)
SMD (95% CI)		-0.4 (-0.7 to -0.1)			

(continued)

Table 2. Mean Symptom Scores at the End of Therapy According to Treatment (continued)

Source	Outcome Assessment Tool ^a	Active Treatment		Placebo	
		No. of Participants	Score, Mean (SD)	No. of Participants	Score, Mean (SD)
Quality-of-Life Score					
α-Blockers					
Nickel et al, ⁹ 2008	NIH-CPSI	138	3.3 (2.5)	134	3.5 (2.3)
Tuğcu et al, ¹⁰ 2007	NIH-CPSI	30	3.8 (1.1)	30	6.9 (1.1)
Alexander et al, ²¹ 2004	NIH-CPSI	45	6.9 (3.4)	45	7.1 (3.3)
Cheah et al, ³³ 2003	NIH-CPSI	43	3.6 (3.4)	43	5.5 (3.9)
Eviyaoğlu and Burgut, ³⁴ 2002	IPSS	30	2.3 (0.8)	30	3.7 (0.8)
SMD (95% CI)		-1.0 (-1.8 to -0.2)			
Antibiotics					
Alexander et al, ²¹ 2004	NIH-CPSI	42	5.8 (3.9)	45	7.1 (3.3)
Nickel et al, ³⁹ 2003	NIH-CPSI	45	3.7 (1.8)	35	3.8 (1.7)
Zhou et al, ⁴⁴ 2008	NIH-CPSI	24	5.5 (0.6)	24	8.5 (1.0)
USMD (95% CI)		-1.5 (-3.6 to 0.6)			

Abbreviations: CI, confidence interval; SMD, standardized mean difference; USMD, unstandardized mean difference.

^aInternational Prostate Symptom Score (IPSS) ranges from 0 to 51; National Institutes of Health Chronic Prostatitis Symptom Index (NIH-CPSI) ranges from 0 to 43; Prostatitis Symptom Score Index (PSSI) ranges from 0 to 12; and pain and voiding questionnaires range from 0 to 3.

dom-effects model was -1.1 (95% CI, -1.8 to -0.3). Patients receiving α-blockers had a 1.1-unit (equivalent to 2.2 points for the NIH-CPSI or 1.9 points for the PSSI/pain questionnaire) significantly lower pain score than patients who received placebo, but this was highly heterogeneous across studies (*I*²=94%). The source of this heterogeneity was not apparent. A sensitivity analysis limited to the 4 studies^{9,21,33,34} with adequate sequence generation yielded an SMD of -0.3 (95% CI, -0.5 to -0.1). The Egger test suggested publication bias due to small-study effect (coefficient=-8.9; *P*=.02) (eFigure 2B). Adjusting for this bias removed the beneficial effect of α-blockers (coefficient=1.2; 95% CI, -0.1 to 2.4; *P*=.06).

Three studies^{21,39,44} with a total of 215 patients compared mean pain scores between antibiotics and placebo. The USMD was -2.7 (95% CI, -8.7 to 3.2); ie, the mean pain score in the antibiotic group was 2.7 NIH-CPSI units lower than in the placebo group, a difference that was not statistically significant (Table 2). There was no evidence of publication bias (coefficient=-9.3; SE, 4.2; *P*=.27).

Three studies¹³⁻¹⁵ comparing phytotherapy and placebo were pooled (n=222). The SMD was -0.5 (95% CI, -0.7 to -0.2); ie, patients receiving phytotherapy had a 0.5-unit significantly lower pain score than patients receiving placebo (Table 2). There was no evidence of publication bias (coefficient=-2.0; *P*=.08).

Voiding Scores. Five studies^{9,10,21,33,34} compared α-blockers and placebo for the outcome of voiding scores (n=568). In the random-effects model, the SMD was -1.4 (95% CI, -2.3 to -0.5). That is, the mean voiding scores were 1.4 units significantly lower in the α-blocker groups than in the placebo groups (Table 2 and eFigure 1C). This difference translates into 3.1 units on the NIH-CPSI and IPSS scales. In sensitivity analyses limited to the 4 studies^{9,21,33,34} with adequate randomization, the SMD was -0.7 (95% CI, -1.2 to -0.2). The Egger test suggested publication bias due to small-study effects (coefficient=-9.5; *P*=.02). The contour-enhanced funnel plot indicated asymmetry (eFigure 2C), and adjusting for this bias removed any favorable treatment effect (coefficient=1.1; 95% CI, -0.4 to 2.6; *P*=.10).

Three studies^{21,39,44} comparing antibiotics and placebo (n=215) had a pooled SMD of -3.2 (95% CI, -6.1 to -0.3) (Table 2). Three studies comparing phytotherapy with placebo (n=223)¹³⁻¹⁵ had an SMD of -0.4 (95% CI, -0.7 to -0.1) (Table 2).

Quality-of-Life Scores. Five studies compared quality-of-life scores between α-blocker and placebo groups (n=568).^{9,10,21,33,34} The SMD was -1.0 (95% CI, -1.8 to -0.2) (or approximately 1.4 units on the NIH-CPSI and IPSS scales) lower in the α-blocker group than in the placebo group. How-

ever, results were heterogeneous (eFigure 1D). The contour-enhanced funnel plot suggested publication bias from 2 small studies^{10,34} that had strong treatment effects (eFigure 2D); adjusting for this bias removed the significant benefit of α-blockers (coefficient=1.2; 95% CI, -0.6 to 2.9; *P*=.13).

Three studies^{21,39,44} (n=215) compared quality of life between antibiotics and placebo. The estimated USMD was -1.5 (95% CI, -3.6 to 0.6) NIH-CPSI units lower in antibiotics groups than placebo groups, but this difference was not statistically significant (Table 2). The Egger test did not suggest publication bias (coefficient=-13.2; SE, 2.7; *P*=.13).

Pooled Response Rate. Nine studies[†] included a dichotomized treatment response as the outcome of interest. Of these, 6 studies^{9,10,21,24,33,38} compared α-blockers with placebo and 3 studies^{12,22,43} compared anti-inflammatory drugs with placebo. Among 6 studies (n=602) comparing α-blockers with placebo, various criteria were used for assessing treatment responses (TABLE 3). The pooled RR was 1.6 (95% CI, 1.1-2.3); ie, patients receiving α-blockers were 60% more likely to have a response than patients receiving placebo (eFigure 3A). However, there was moderately high heterogeneity. Meta-regression evaluating duration of treatment reduced the *I*² from 64.2% to 12.8%, indicating that

[†]References 9, 10, 12, 21, 22, 24, 33, 38, 43.

Table 3. Treatment Response Rates for α -Blockers and Anti-inflammatory Drugs

Source	Definition of Treatment Response	Active Treatment		Placebo		RR (95% CI)
		No. of Responses	No. of Nonresponses	No. of Responses	No. of Nonresponses	
α-Blockers						
Nickel et al, ⁹ 2008	4-point decrease in NIH-CPSI	68	70	66	68	1.0 (0.8-1.3)
Tuğcu et al, ¹⁰ 2007	50% decrease in NIH-CPSI	20	10	10	20	2.0 (1.4-3.5)
Alexander et al, ²¹ 2004	4-point decrease in NIH-CPSI	12	33	11	34	1.1 (0.5-2.3)
Nickel et al, ²⁴ 2004	50% decrease in NIH-CPSI	9	18	5	25	2.0 (0.8-5.2)
Cheah et al, ³³ 2003	50% decrease in NIH-CPSI	24	19	14	29	1.6 (1.0-2.6)
Mehik et al, ³⁸ 2003	33% decrease in NIH-CPSI	13	4	4	16	2.5 (1.4-4.5)
Pooled RR						1.6 (1.1-2.3)
Anti-inflammatory drugs						
Nickel et al, ¹² 2003	25% decrease in NIH-CPSI	31	18	24	35	1.6 (1.1-2.6)
Bates et al, ²² 2007	6-point decrease in NIH-CPSI	2	4	4	8	1.0 (0.3-4.0)
Zhao et al, ⁴³ 2009	25% decrease in NIH-CPSI	25	7	10	22	2.5 (1.5-4.3)
Pooled RR						1.8 (1.2, 2.6)

Abbreviations: CI, confidence interval; NIH-CPSI National Institutes of Health Chronic Prostatitis Symptom Index; RR, relative risk.

treatment duration may explain the heterogeneity (eFigure 3A). Duration of treatment was 6 to 12 weeks in 3 studies^{9,21,24} and 14 to 24 weeks in the other 3 studies.^{10,33,38} Pooled RRs within these subgroups were homogeneous (eFigure 3A) at 1.0 (95% CI, 0.8-1.3) for 6 to 12 weeks' duration and 2.0 (95% CI, 1.4-3.0) for 14 to 24 weeks' duration, respectively.

The contour-enhanced funnel plot suggested asymmetry, especially within the 3 studies in which treatment duration ranged from 14 to 24 weeks (eFigure 3B). Metatrim indicated that 3 studies with negative effects of α -blockers were missing. Adding these studies into pooling yielded no benefit from α -blockers, with a pooled RR of 1.1 (95% CI, 0.8-1.7).

Three studies^{12,22,43} were pooled to compare anti-inflammatory therapies with placebo (n=190). The types of anti-inflammatory drugs were cyclooxygenase 2 inhibitors in 2 studies^{12,43} and a corticosteroid in 1 study.²² The pooled RR was 1.8 (95% CI, 1.2-2.6), with mild heterogeneity ($I^2=24.4%$). These results indicate that patients receiving anti-inflammatory drugs were 80% more likely to have a favorable response than patients receiving placebo (Table 3). The Egger test did not suggest publication bias (coefficient=-0.36; $P=.90$).

Network Meta-analysis

Total Symptom Scores. Data from 13 studies (n=1541)^{9,10,14,15,21,33,39-45} using

the NIH-CPSI were included in network meta-analyses for the outcome of total symptom score (eTable 2). Treatment comparisons are described in eTable 3 and eFigure 4. Mean total scores at follow-up were, for α -blockers, -11.0 (95% CI, -13.9 to -8.1), for antibiotics, -9.8 (95% CI, -15.1 to -4.6), for α -blockers plus antibiotics, -13.8 (95% CI, -17.5 to -10.2), and for finasteride, -4.6 (95% CI, -8.7 to -0.5) units significantly lower than placebo groups. In this instance, α -blockers plus antibiotics were better than any other therapy and were significantly better than α -blockers alone (-2.9; 95% CI, -5.2 to -0.5).

Pain Scores. The network meta-analysis was performed with 14 studies[‡] that used similar NIH-CPSI scores (eTable 2). α -Blockers, antibiotics, α -blockers plus antibiotics, and anti-inflammatory drugs were associated with significantly better pain scores at follow-up than placebo, with pain scores of -4.1 (95% CI, -5.9 to -2.3), -4.4 (95% CI, -7.0 to -1.9), -5.7 (95% CI, -7.8 to -3.6), and -3.0 (95% CI, -5.7 to -0.4), respectively (eTable 3 and eFigure 5). Again, the most favorable therapy was α -blockers plus antibiotics (eFigure 5).

Voiding Scores. Thirteen studies[§] (n=631) were included in the voiding analysis (eTable 2 and eFigure 6). Mean voiding scores at follow-up for α -blockers, antibiotics, and α -blockers plus

antibiotics were -3.4 (95% CI, -4.8 to -2.1), -2.8 (95% CI, -4.1 to -1.6), and -3.7 (95% CI, -5.2 to -2.1) units significantly lower than for placebo (eTable 3). Again, the best treatment in the network comparisons was α -blockers plus antibiotics.

Quality-of-Life Scores. Twelve studies (n=1477)^{||} were included in the quality-of-life analysis (eTable 2). Associations of α -blockers, antibiotics, and α -blockers plus antibiotics (-1.9; 95% CI, -3.6 to -0.2) were significantly better than placebo (eTable 3 and eFigure 7). α -Blockers plus antibiotics was the best treatment in the network comparisons, with a decrease of -2.8 (95% CI, -4.7 to -0.9) units in the quality-of-life score.

Response Rate. Fourteen studies[¶] (n=1561) reported favorable response to treatment (eFigure 8 and eTable 4). The RR of treatment response was highest with anti-inflammatory drugs (RR, 1.8; 95% CI, 1.3-2.6), followed by phytotherapy (RR, 1.6; 95% CI, 1.1-2.4) and α -blockers (RR, 1.3; 95% CI, 1.0-1.6) compared with placebo (eTable 5). Anti-inflammatory therapies were the best treatment in the network comparisons.

COMMENT

We performed a systematic review and meta-analysis of outpatient treatments for CP/CPPS. We studied relevant clinical

‡References 9-11, 13-15, 21, 33, 39, 41-45.
§References 9, 10, 13-15, 21, 33, 39, 41-45.

||References 9, 10, 14, 15, 21, 33, 39, 41-45.
¶References 9, 10, 12, 15, 21, 22, 24, 33, 36, 38-40, 43, 45.

outcomes, including total clinical symptom, voiding, pain, and quality-of-life scores, as well as treatment response rates. α -Blockers, antibiotics, and a combination of the 2 appear to improve all clinical symptom scores compared with placebo, while anti-inflammatory drugs, finasteride, and phytotherapies have a lesser but measurable effect on select variables (ie, pain, voiding symptoms, and treatment response rate, respectively). However, the treatment effects of α -blockers are distorted by publication bias/small-study effects, and adjusting for this removes any treatment benefit.

Given the large number of treatment options, meta-analyses of direct comparisons are limited by the small number of studies that evaluated a particular pair of treatments. Network meta-analysis circumvents this problem by creating indirect comparisons and allowing data synthesis that can help identify the most effective therapy. In this case, α -blockers plus antibiotics was consistently the best therapy when the outcome was symptom scores. Anti-inflammatory drugs were the best therapy when treatment response was the outcome; although steroids and nonsteroidal anti-inflammatory drugs were pooled together because of the small number of studies, the heterogeneity was low, indicating that their effects may be similar.

Treatment benefits (whether we measured effect on symptom scores or responder rates) were modest for some therapies and nonexistent for others. This probably reflects the heterogeneous nature of patients presenting with CP/CPPS. Patients with CP/CPPS represent a group of divergent clinical phenotypes based on the various etiologies and pathogenic pathways that underlie this enigmatic condition.⁴⁶ As a result of difficulties in diagnoses, some patients (in clinical trials and practice) likely receive inappropriate therapies. It makes clinical sense that patients with predominant voiding dysfunction may respond best to α -blockers, those with a history of urinary tract infection may respond best to antibiotics, and those with pain/inflammation may respond best to anti-inflammatory drugs and/or gabapenti-

noids. However, further study is needed to determine whether patient characteristics determine the most effective therapy for CP/CPPS.

Because the diagnosis of CP/CPPS requires exclusion of infection, the reason for the benefit associated with antibiotics is not immediately clear. This observed effect may be due to the eradication or suppression of uncultured or unrecognized uropathogens that may be measurable with polymerase chain reaction.⁴⁷⁻⁴⁹ In addition, it is important to point out that quinolones have anti-inflammatory^{50,51} and analgesic properties.⁵²

While the results of our analyses demonstrate that α -blockers, antibiotics, and anti-inflammatory medications are beneficial for CP/CPPS, we recognize that the total sample sizes are relatively small and that the effect sizes are modest and often below the minimal clinically significant difference. Furthermore, even these estimates may be overinflated given the evidence for publication bias. Our results suggest that future research should focus on using these treatments rationally, perhaps using individualized patient therapy or multiple therapies directed toward the specific clinical phenotype of each unique patient.⁵³ This concept is presently being evaluated.⁴⁹ The decision to use these therapies also needs to take account of the adverse event profile: α -blockers can cause postural hypotension, edema, and drowsiness; quinolones can cause dizziness, headaches, and gastrointestinal upset; and nonsteroidal anti-inflammatory drugs can cause gastritis, renal impairment, and edema. The risk-benefit ratio remains to be determined.

Our study has several strengths. The review methods were systematic and exhaustive. Contour-enhanced funnel plots helped to identify possible publication bias due to small-study effects, which tend to lead to higher treatment effects than large studies.^{25,26} We mapped all possible treatment comparisons (9 treatments with 36 possible pairwise comparisons) using a network meta-analysis.²⁷⁻²⁹ An advantage of network meta-analysis is the ability to "borrow" information on the treatment

groups from other studies, thereby increasing the total sample sizes. For example, direct comparisons in a previous meta-analysis¹⁶ included 466, 236, and 123 patients in pooling for total symptom, pain, and voiding scores, respectively, comparing α -blockers vs placebo. In the current analysis, the corresponding numbers of patients were 1549, 1556, and 1546. We applied a mixed model, which is thought to be the most appropriate method for network meta-analysis.^{27,54} Although our pooled estimates were quite heterogeneous, the mixed model with random intercept takes into account variations at the study level.

These methods have limitations, however. Although all studies were randomized controlled trials, most studies were unclear in randomization sequence generations and, hence, selection bias or confounding might be present. Pooled results were often heterogeneous and the source of this difference was not apparent.

CONCLUSION

Our review suggests that α -blockers, antibiotics, or combinations of both are most appropriate for therapy of CP/CPPS, particularly for patients with voiding symptoms. However, the magnitude of apparent benefit with α -blockers may be distorted by publication bias. Anti-inflammatory medications remain an option for patients presenting with pain. While finasteride and phytotherapy may provide benefit to some patients, these therapies require more evaluation, perhaps in selected subgroups of CP/CPPS patients.

Author Affiliations: Section for Clinical Epidemiology and Biostatistics (Drs Anothaisintawee, Numthavaj, and Thakkinian) and Department of Family Medicine (Drs Anothaisintawee and Thammakraisorn), Faculty of Medicine, Ramathibodi Hospital, Mahidol University, Bangkok, Thailand; Centre for Clinical Epidemiology and Biostatistics, School of Medicine and Public Health, University of Newcastle (Dr Attia and Mr McEvoy), and Hunter Medical Research Institute (Dr Attia), Newcastle, New South Wales, Australia; and Department of Urology, Queens University, Kingston, Ontario, Canada (Dr Nickel).

Author Contributions: Dr Thakkinian had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Thakkinian.

Acquisition of data: Anothaisintawee, Thammakraisorn, Numthavaj, Thakkinian.

Analysis and interpretation of data: Anothaisintawee, Attia, Nickel, McEvoy, Thakkinian.

Drafting of the manuscript: Anothaisintawee, Thakkinstian.

Critical revision of the manuscript for important intellectual content: Attia, Nickel, Thammakraisorn, Numthavaj, McEvoy, Thakkinstian.

Statistical analysis: Anothaisintawee, Thakkinstian.

Administrative, technical, or material support: Thammakraisorn, Numthavaj, Thakkinstian.

Study supervision: Attia, Nickel, Thakkinstian.

Conflict of Interest Disclosures: All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Dr Nickel reported serving as a consultant for GlaxoSmithKline, Pfizer, Watson, Farr Labs, Astellas, and Triton and as an investigator for GlaxoSmithKline, Pfizer, and Watson. No other authors reported disclosures.

Funding/Support: Dr Nickel's research in CP/CPPS is funded in part by the National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, the Canada Institute for Health Research, and the Canada Research Chair Program.

Role of the Sponsor: None of the funding organizations or sponsors had any role in the design and conduct of the study; the extraction, management, analysis, or interpretation of the data; or the preparation, review, or approval of the manuscript.

Online-Only Materials: The eAppendix, eTables 1 through 6, and eFigures 1 through 8 are available online at <http://www.jama.com>.

REFERENCE

- Roberts RO, Lieber MM, Rhodes T, et al. Prevalence of a physician-assigned diagnosis of prostatitis. *Urology*. 1998;51(4):578-584.
- Collins MM, Stafford RS, O'Leary MP, Barry MJ. How common is prostatitis? *J Urol*. 1998;159(4):1224-1228.
- de la Rosette JJ, Hubregtse MR, Meuleman EJ, et al. Diagnosis and treatment of 409 patients with prostatitis syndromes. *Urology*. 1993;41(4):301-307.
- Krieger JN, Nyberg L Jr, Nickel JC. NIH consensus definition and classification of prostatitis. *JAMA*. 1999;282(3):236-237.
- McNaughton Collins M, Pontari MA, O'Leary MP, et al. Chronic Prostatitis Collaborative Research Network. Quality of life is impaired in men with chronic prostatitis. *J Gen Intern Med*. 2001;16(10):656-662.
- McNaughton Collins M, MacDonald R, Wilt TJ. Diagnosis and treatment of chronic abacterial prostatitis. *Ann Intern Med*. 2000;133(5):367-381.
- Hetrick DC, Ciol MA, Rothman I, et al. Musculoskeletal dysfunction in men with chronic pelvic pain syndrome type III. *J Urol*. 2003;170(3):828-831.
- Hochreiter WW, Nadler RB, Koch AE, et al. Evaluation of the cytokines interleukin 8 and epithelial neutrophil activating peptide 78 as indicators of inflammation in prostatic secretions. *Urology*. 2000;56(6):1025-1029.
- Nickel JC, Krieger JN, McNaughton-Collins M, et al. Chronic Prostatitis Collaborative Research Network. Alfuzosin and symptoms of chronic prostatitis-chronic pelvic pain syndrome. *N Engl J Med*. 2008;359(25):2663-2673.
- Tuğcu V, Taşçı AI, Fazlıoğlu A, et al. A placebo-controlled comparison of the efficiency of triple- and monotherapy in category III B chronic pelvic pain syndrome (CPPS). *Eur Urol*. 2007;51(4):1113-1117.
- Goldmeier D, Madden P, McKenna M, Tamm N. Treatment of category III A prostatitis with zafirlukast: a randomized controlled feasibility study. *Int J STD AIDS*. 2005;16(3):196-200.
- Nickel JC, Pontari M, Moon T, et al. Rofecoxib Prostatitis Investigator Team. A randomized, placebo controlled, multicenter study to evaluate the safety and efficacy of rofecoxib in the treatment of chronic non-bacterial prostatitis. *J Urol*. 2003;169(4):1401-1405.
- Elist J. Effects of pollen extract preparation Prostat/Polit on lower urinary tract symptoms in patients with chronic nonbacterial prostatitis/chronic pelvic pain syndrome: a randomized, double-blind, placebo-controlled study. *Urology*. 2006;67(1):60-63.
- Shoskes DA, Zeitlin SI, Shahed A, Rajfer J. Quercetin in men with category III chronic prostatitis. *Urology*. 1999;54(6):960-963.
- Wagenlehner FM, Schneider H, Ludwig M, et al. A pollen extract (Cernilton) in patients with inflammatory chronic prostatitis-chronic pelvic pain syndrome. *Eur Urol*. 2009;56(3):544-551.
- Yang G, Wei Q, Li H, et al. The effect of α -adrenergic antagonists in chronic prostatitis/chronic pelvic pain syndrome. *J Androl*. 2006;27(6):847-852.
- Higgins JPT, Altman DG. Assessing risk of bias in included studies. In: Higgins JPT, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions*. Chichester, England: John Wiley & Sons; 2008.
- Litwin MS, McNaughton-Collins M, Fowler FJ Jr, et al. Chronic Prostatitis Collaborative Research Network. The National Institutes of Health Chronic Prostatitis Symptom Index. *J Urol*. 1999;162(2):369-375.
- Cockett ATK, Khoury S, Aso Y, et al. *Second International Consultation on Benign Prostatic Hyperplasia*. Hong Kong: Scientific Communications International; 1994:624-635.
- Krieger JN, Egan KJ, Ross SO, et al. Chronic pelvic pains represent the most prominent urogenital symptoms of "chronic prostatitis." *Urology*. 1996;48(5):715-721.
- Alexander RB, Proppert KJ, Schaeffer AJ, et al. Chronic Prostatitis Collaborative Research Network. Ciprofloxacin or tamsulosin in men with chronic prostatitis/chronic pelvic pain syndrome. *Ann Intern Med*. 2004;141(8):581-589.
- Bates SM, Hill VA, Anderson JB, et al. A prospective, randomized, double-blind trial to evaluate the role of a short reducing course of oral corticosteroid therapy in the treatment of chronic prostatitis/chronic pelvic pain syndrome. *BJU Int*. 2007;99(2):355-359.
- Nickel JC. Treatment of chronic prostatitis/chronic pelvic pain syndrome. *Int J Antimicrob Agents*. 2008;31(suppl 1):S112-S116.
- Nickel JC, Narayan P, McKay J, Doyle C. Treatment of chronic prostatitis/chronic pelvic pain syndrome with tamsulosin. *J Urol*. 2004;171(4):1594-1597.
- Moreno SG, Sutton AJ, Turner EH, et al. Novel methods to deal with publication biases. *BMJ*. 2009;339:b2981.
- Nüesch E, Trelle S, Reichenbach S, et al. Small study effects in meta-analyses of osteoarthritis trials. *BMJ*. 2010;341:c3515.
- Lu G, Ades AE. Combination of direct and indirect evidence in mixed treatment comparisons. *Stat Med*. 2004;23(20):3105-3124.
- Song F, Altman DG, Glenny AM, Deeks JJ. Validity of indirect comparison for estimating efficacy of competing interventions. *BMJ*. 2003;326(7387):472.
- Song F, Harvey I, Lilford R. Adjusted indirect comparison may be less biased than direct comparison for evaluating new pharmaceutical interventions. *J Clin Epidemiol*. 2008;61(5):455-463.
- Stata [computer program]. Release 11. College Station, TX: Stata Corp; 2009.
- Kaplan SA, Volpe MA, Te AE. A prospective, 1-year trial using saw palmetto vs finasteride in the treatment of category III prostatitis/chronic pelvic pain syndrome. *J Urol*. 2004;171(1):284-288.
- Kulovac B, Aganović D, Prcić A, Hadzirosmanović O. Management of chronic nonbacterial prostatitis/chronic pelvic pain syndrome. *Bosn J Basic Med Sci*. 2007;7(3):245-249.
- Cheah PY, Liong ML, Yuen KH, et al. Terazosin therapy for chronic prostatitis/chronic pelvic pain syndrome. *J Urol*. 2003;169(2):592-596.
- Evlıyaoğlu Y, Burgut R. Lower urinary tract symptoms, pain and quality of life assessment in chronic non-bacterial prostatitis patients treated with α -blocking agent doxazosin; vs placebo. *Int Urol Nephrol*. 2002;34(3):351-356.
- Gül O, Eroğlu M, Ozok U. Use of terazosin in patients with chronic pelvic pain syndrome and evaluation by Prostatitis Symptom Score Index. *Int Urol Nephrol*. 2001;32(3):433-436.
- Jeong CW, Lim DJ, Son H, et al. Treatment for chronic prostatitis/chronic pelvic pain syndrome: levofloxacin, doxazosin and their combination. *Urol Int*. 2008;80(2):157-161.
- Leskinen M, Lukkarinen O, Marttila T. Effects of finasteride in patients with inflammatory chronic pelvic pain syndrome. *Urology*. 1999;53(3):502-505.
- Mehik A, Alas P, Nickel JC, et al. Alfuzosin treatment for chronic prostatitis/chronic pelvic pain syndrome. *Urology*. 2003;62(3):425-429.
- Nickel JC, Downey J, Clark J, et al. Levofloxacin for chronic prostatitis/chronic pelvic pain syndrome in men. *Urology*. 2003;62(4):614-617.
- Nickel JC, Downey J, Pontari MA, et al. A randomized placebo-controlled multicentre study to evaluate the safety and efficacy of finasteride for male chronic pelvic pain syndrome (category IIIA chronic nonbacterial prostatitis). *BJU Int*. 2004;93(7):991-995.
- Nickel JC, Forrest JB, Tomera K, et al. Pentosan polysulfate sodium therapy for men with chronic pelvic pain syndrome. *J Urol*. 2005;173(4):1252-1255.
- Ye ZQ, Lan RZ, Yang WM, et al. Tamsulosin treatment of chronic non-bacterial prostatitis. *J Int Med Res*. 2008;36(2):244-252.
- Zhao WP, Zhang ZG, Li XD, et al. Celecoxib reduces symptoms in men with difficult chronic pelvic pain syndrome (category IIIA). *Braz J Med Biol Res*. 2009;42(10):963-967.
- Zhou Z, Hong L, Shen X, et al. Detection of nanobacteria infection in type III prostatitis. *Urology*. 2008;71(6):1091-1095.
- Pontari MA, Krieger JN, Litwin MS, et al. Chronic Prostatitis Collaborative Research Network-2. Pregabalin for the treatment of men with chronic prostatitis/chronic pelvic pain syndrome. *Arch Intern Med*. 2010;170(17):1586-1593.
- Shoskes DA, Nickel JC, Dolinga R, Prots D. Clinical phenotyping of patients with chronic prostatitis/chronic pelvic pain syndrome and correlation with symptom severity. *Urology*. 2009;73(3):538-542.
- Nickel JC, Moon T. Chronic bacterial prostatitis. *Urology*. 2005;66(1):2-8.
- Nickel JC, Xiang J. Clinical significance of non-traditional bacterial uropathogens in the management of chronic prostatitis. *J Urol*. 2008;179(4):1391-1395.
- Shoskes DA, Nickel JC, Kattan MW. Phenotypically directed multimodal therapy for chronic prostatitis/chronic pelvic pain syndrome. *Urology*. 2010;75(6):1249-1253.
- Galley HF, Nelson SJ, Dubbels AM, Webster NR. Effect of ciprofloxacin on the accumulation of interleukin-6, interleukin-8, and nitrite from a human endothelial cell model of sepsis. *Crit Care Med*. 1997;25(8):1392-1395.
- Yoshimura T, Kurita C, Usami E, et al. Immunomodulatory action of levofloxacin on cytokine production by human peripheral blood mononuclear cells. *Chemotherapy*. 1996;42(6):459-464.
- Katsuno G, Takahashi HK, Iwagaki H, et al. The immunosuppressive effects of ciprofloxacin during human mixed lymphocyte reaction. *Clin Immunol*. 2006;119(1):110-119.
- Nickel JC, Shoskes DA. Phenotypic approach to the management of the chronic prostatitis/chronic pelvic pain syndrome. *BJU Int*. 2010;106(9):1252-1263.
- Caldwell DM, Ades AE, Higgins JP. Simultaneous comparison of multiple treatments. *BMJ*. 2005;331(7521):897-900.