

Case 4

A 2-year-old girl from Nakornrachasima

Chief complaint multiple flaccid bullae since birth

Present illness At birth she had a flaccid bullae sized 6 cm on her right elbow. The lesions progress in number especially on pressure and traumatic area. There are serosanguinous fluid in the bullae. The lesions turn scar when they rupture.

Past history She is a first child in her family, She is delivered by vacuum extraction. Her body weight is 3,000 grams. She has normal growth and development

Family history : Her Great grandmother, Grandfather, Father and two uncles also have the same lesions. (shown in pedigree) Their lesions are all resolved leaving hypopigmented scars, except for one uncle that still has blistering lesions.

Physical examination A Thai young girl, alert, not pale, no jaundice

HEENT : pharynx & tonsils are not injected, not pale conjunctiva,
no cervical lymphadenopathy

Heart : normal S1S2, no murmur

Lung : clear, no adventitious sound

Abdomen : soft, no hepatosplenomegaly

Skin : flaccid bullae with clear fluid and serosanguinous fluid at right foot

: multiple hypopigmented scarring patch and macules at trunk and extremities



Fig.1

Histopathology pending
EM: pending

Diagnosis Epidermolysis Bullosa
Presenter Sutasinee Tunsuriyawong
Consultant Wasuthida Pabunrueng